



Generating the Pfaffian closure with total Pfaffian functions

GARETH JONES
PATRICK SPEISSEGGER

Abstract: Given an o-minimal expansion \mathcal{R} of the real field, we show that the structure obtained from \mathcal{R} by iterating the operation of adding all total Pfaffian functions over \mathcal{R} defines the same sets as the Pfaffian closure of \mathcal{R} .

2010 Mathematics Subject Classification [14P10,03C64](#) (primary); [58A17](#) (secondary)

Keywords: o-minimal structure, Pfaffian function, Rolle leaf

There are various possibilities for adding Pfaffian objects to o-minimal expansions of the real field and preserving o-minimality. One example is the Pfaffian closure of an o-minimal expansion of the real field, which was shown to be o-minimal by the second author [7]. The purpose of this note is to present a somewhat simpler construction of the Pfaffian closure. Although not as simple as the description in terms of nested leaves obtained by Lion and the second author [5], our construction has the novelty of only using total Pfaffian functions and is reminiscent of the original Pfaffian expansion of the real field constructed by Wilkie [9].

In order to state our result, we need to introduce some terminology. Suppose that \mathcal{R} is an o-minimal expansion of the real field, and that $U \subseteq \mathbb{R}^n$ is an \mathcal{R} -definable open subset of \mathbb{R}^n for some $n \in \mathbb{N}$. We say that a C^1 function $f : U \rightarrow \mathbb{R}$ is *Pfaffian over \mathcal{R}* if there exist \mathcal{R} -definable C^1 functions $P_i : U \times \mathbb{R} \rightarrow \mathbb{R}$, for $i = 1, \dots, n$ such that

$$\frac{\partial f}{\partial x_i}(x) = P_i(x, f(x))$$

for all $x \in U$.

Given $n, l \in \mathbb{N}$ such that $l \leq n$, we let G_n^l be the Grassmannian of all linear subspaces of \mathbb{R}^n of dimension l . This is an analytic manifold and is naturally definable in the real field (see [1, 3.4.2]). We also set $G_n = \bigcup_{l=0}^n G_n^l$. Now fix an embedded C^1 submanifold M of \mathbb{R}^n and let $l \leq n$. A C^1 map $d : M \rightarrow G_n$ is said to be a *distribution on M* if $d(x) \subseteq T_x M$ for all $x \in M$, where $T_x M$ is the tangent space of M at x . A distribution d is an *l -distribution* if $d(M) \subseteq G_n^l$. Given an l -distribution d on M and an immersed C^1 submanifold V of M , we say that V is an *integral manifold* of d if $T_x V = d(x)$ for all

$x \in V$. A maximal connected integral manifold is called *leaf* of the distribution. Now suppose that d has codimension one. A leaf L of d is said to be a *Rolle leaf* of d if it is a closed embedded submanifold of M and is such that for all C^1 curves $\gamma : [0, 1] \rightarrow M$ satisfying $\gamma(0), \gamma(1) \in L$, we have $\gamma'(t) \in d(\gamma(t))$ for some $t \in [0, 1]$. A *Rolle leaf over \mathcal{R}* is a Rolle leaf of an \mathcal{R} -definable codimension one distribution defined on \mathbb{R}^n for some $n \in \mathbb{N}$. For example, a result due to Khovanskii (see [8, 1.6]) implies that if $f : \mathbb{R}^n \rightarrow \mathbb{R}$ is Pfaffian over \mathcal{R} , then the graph of f is a Rolle leaf over \mathcal{R} .

We can now define the Pfaffian structures involved in our result. Given any o-minimal expansion of the real field \mathcal{R} , let $\mathcal{L}(\mathcal{R})$ be the collection of all Rolle leaves over \mathcal{R} . Now let $\mathcal{R}_0 = \mathcal{R}$ and, for $i \geq 0$, let \mathcal{R}_{i+1} be the expansion of \mathcal{R}_i by all leaves in $\mathcal{L}(\mathcal{R}_i)$. Let \mathcal{L} be the union of all the $\mathcal{L}(\mathcal{R}_i)$ and let $\mathcal{P}(\mathcal{R})$ be the expansion of \mathcal{R} by all the leaves in \mathcal{L} . This structure is called the *Pfaffian closure* of \mathcal{R} . The second author showed that it is o-minimal [7].

Similarly, we let $\mathcal{L}'(\mathcal{R})$ be the collection of all functions $f : \mathbb{R}^n \rightarrow \mathbb{R}$, for all $n \in \mathbb{N}$ that are Pfaffian over \mathcal{R} . We define \mathcal{R}'_i and then $\mathcal{P}'(\mathcal{R})$ by mimicking the previous paragraph. The structure $\mathcal{P}'(\mathcal{R})$ is a reduct of $\mathcal{P}(\mathcal{R})$ (by the example above) and it is the purpose of this note to show that they are in fact the same from the point of view of definability.

Theorem 1 *A set $X \subseteq \mathbb{R}^n$ is definable in $\mathcal{P}(\mathcal{R})$ if and only if it is definable in $\mathcal{P}'(\mathcal{R})$.*

If \mathcal{R} admits analytic cell decomposition, then so too does $\mathcal{P}'(\mathcal{R})$ (see [8]) and it follows that in this case, the reduct of $\mathcal{P}'(\mathcal{R})$ in which only analytic functions are added also defines the same sets as $\mathcal{P}(\mathcal{R})$.

Before proving the theorem, we first recall a result from Khovanskii theory that we will repeatedly use in the proof. This originates with Khovanskii's work on pfaffian functions (see [3]) and was adapted to the o-minimal setting by the second author [7] following work of Moussu and Roche [6] and Lion and Rolin [4]. In our proof we shall only ever need to work with a single distribution so we restrict ourselves to Khovanskii theory in this simple setting. We follow the presentation in [8]. Suppose that d is a C^2 distribution on a C^2 manifold $M \subseteq \mathbb{R}^n$ and that $N \subseteq M$ is a C^2 submanifold of M . The *pull-back* of d to N is the distribution d^N on N defined by

$$d^N(x) = T_x N \cap d(x).$$

We say that N is compatible with d if d^N is an l -distribution on N , for some l . From now on, we use the word definable to mean $\mathcal{P}'(\mathcal{R})$ -definable. In particular, cell means $\mathcal{P}'(\mathcal{R})$ -definable cell. The result we need is as follows (see [8, 3.6]).

Fact 2 Let d be a definable codimension one distribution on M and suppose that \mathcal{C} is a partition of M into C^2 cells compatible with d . Then there is a $k \in \mathbb{N}$ such that whenever $C \in \mathcal{C}$ and L is a Rolle leaf of d the set $C \cap L$ is a union of at most k Rolle leaves of d^C .

Given the definition of $\mathcal{P}'(\mathcal{R})$, in order to prove the theorem it suffices to show that if L is a Rolle leaf over $\mathcal{P}'(\mathcal{R})$ then L is definable. For the proof of this, we assume that the reader is familiar with o-minimality (as presented in [2]). First, an easy observation.

Lemma 3 Suppose that $C \subseteq \mathbb{R}^n$ is an open C^2 cell and that $f : C \rightarrow \mathbb{R}$ is Pfaffian over $\mathcal{P}'(\mathcal{R})$. Then f is definable.

The proof, using a definable diffeomorphism between C and \mathbb{R}^n , is left to the reader.

Now suppose that $C \subseteq \mathbb{R}^n$ is a bounded open C^2 cell, and that $\alpha, \beta, \gamma, \delta : C \rightarrow \mathbb{R}$ are definable bounded C^2 functions such that

$$\gamma(x) < \alpha(x) < \beta(x) < \delta(x)$$

for all $x \in C$. Let $D = (\alpha, \beta)_C$ and $D' = (\gamma, \delta)_C$ and suppose that d' is a definable integrable n -distribution on D' (for a discussion of integrability in this context, see [8, Section 1]). Suppose that we are given a Rolle leaf L' of d' . Assume that both the graph of α and the graph of β are compatible with d' and let d^α and d^β be the pullbacks of d' to the graphs of α and β respectively. Let d be the restriction of d' to C . By Fact 2, $L' \cap D, L' \cap \text{graph}\alpha$ and $L' \cap \text{graph}\beta$ are finite unions of Rolle leaves of d, d^α and d^β respectively.

Lemma 4 Suppose that L is a connected component of $L' \cap D$ and suppose that $\text{graph}\alpha$ is transverse to d' . Then $\text{fr}L \cap \text{graph}\alpha$ is a clopen subset of $L' \cap \text{graph}\alpha$.

Proof Since L' is a Rolle leaf in D' , it is closed in D' and so L is closed in D . So, $\text{fr}L \cap \text{graph}\alpha = \text{cl}L \cap \text{graph}\alpha$ is closed in the graph of α . Using the fact that L' is closed in D' again, we have $\text{cl}L \cap \text{graph}\alpha \subseteq L' \cap \text{graph}\alpha$ and so $\text{fr}L \cap \text{graph}\alpha$ is a closed subset of $L' \cap \text{graph}\alpha$.

We now need to show that $\text{fr}L \cap \text{graph}\alpha$ is open in $L' \cap \text{graph}\alpha$, so let $p \in \text{fr}L \cap \text{graph}\alpha$. Let L_p be the connected component of $L' \cap \text{graph}\alpha$ containing p . By the Frobenius theorem (see [8, Section 1]) there is a neighbourhood U of p and a diffeomorphism $\phi : \mathbb{R}^{n+1} \rightarrow U$ such that $\phi^*d' = \ker dx_{n+1}$ and $\phi(0) = p$. Now, L' is a leaf of d' and $p \in L' \cap U$, so the hyperplane $\mathbb{R}^n \times \{0\}$ is a component of $\phi^{-1}(L' \cap U)$. Since $L' \cap \text{graph}\alpha \cap U$ is a submanifold of $L' \cap U$, we can find an open box B centred at 0

such that $N := \phi^{-1}(L' \cap \text{graph}\alpha \cap U) \cap B$ is connected. Let $B_0 = (\mathbb{R}^n \times \{0\}) \cap B$. Then N is a closed codimension one submanifold of B_0 and so $B_0 \setminus N$ has exactly two components, B_1 and B_2 , say. Since $p \in \text{cl } L$, at least one of B_1 or B_2 must be contained in $\phi^{-1}(L \cap U)$. Also, $N = \text{fr}(B_i) \cap B_0$ for each i and so $\phi(N)$ is contained in $\text{fr}L \cap \text{graph}\alpha$. But $\phi(N)$ is open in $L' \cap \text{graph}\alpha$, by our choice of B , and the lemma is proved. \square

The following proposition suffices to prove the theorem.

Proposition 5 *Let $L \subseteq \mathbb{R}^n$ be a Rolle leaf over $\mathcal{P}'(\mathcal{R})$. Then L is definable in $\mathcal{P}'(\mathcal{R})$.*

Proof The proof is by induction on n . The $n = 1$ case is trivial, so we assume that $n > 1$ and that the proposition is true for Rolle leaves over $\mathcal{P}'(\mathcal{R})$ contained in \mathbb{R}^m with $m < n$. Thus if $C \subseteq \mathbb{R}^n$ is a C^2 cell of dimension less than n and $V \subseteq C$ is a Rolle leaf of a definable codimension one distribution on C , then V is definable.

Suppose that $L \subseteq \mathbb{R}^n$ is a Rolle leaf over $\mathcal{P}'(\mathcal{R})$. Then L is a closed embedded proper submanifold of \mathbb{R}^n , and so there are $p \in \mathbb{R}^n \setminus L$ and $r > 0$ such that $B(p, 2r) \cap L = \emptyset$, where $B(a, \varepsilon)$ is the open ball around a of radius ε . Perhaps after translating and stretching, we may assume that $p = 0$ and that $r = 1$. Let $\phi : \mathbb{R}^n \setminus \{0\} \rightarrow \mathbb{R}^n \setminus \{0\}$ be the semialgebraic diffeomorphism $\phi(x) = \frac{x}{\|x\|^2}$. Then $\phi(L)$ is contained in $B(0, 1/2)$ and $\text{cl}(\phi(L)) \subseteq \phi(L) \cup \{0\}$. So, after replacing L by $\phi(L)$, we may assume that L is a Rolle leaf of a definable integrable $(n - 1)$ -distribution d on $B'(0, 1) := B(0, 1) \setminus \{0\}$, that $L \subseteq B(0, 1/2)$ and that $\text{cl } L \subseteq L \cup \{0\}$.

Let Π_{n-1} be the projection onto the first $n - 1$ coordinates. For each coordinate permutation σ on \mathbb{R}^n , the set $B_\sigma = \{x \in B'(0, 1) : \Pi_{n-1}|_{\sigma^*(d(\sigma^{-1}(x)))}$ has rank $n - 1\}$ is open and together these sets cover $B'(0, 1)$. So it suffices to show that $L \cap B_\sigma$ is definable for each σ . Fix σ , which we may assume to be the identity. Let \mathcal{C} be a C^2 cell decomposition of $B'(0, 1)$ compatible with $B_{\text{id}}, B'(0, 1/2)$ and d . We show that $C \cap L$ is definable for each cell $C \in \mathcal{C}$ such that $C \subseteq B_{\text{id}}$. If $C \in \mathcal{C}$ is not open then $L \cap C$ is definable, by Fact 2 and the inductive hypothesis. So, suppose that $C \in \mathcal{C}$ is open and that $C \subseteq B_{\text{id}}$. Let N be a component of $L \cap C$. Since N is a Rolle leaf of $d|_C$ and C is a cell, N is the graph of a function $f : \Pi_{n-1}(N) \rightarrow \mathbb{R}$. Let $\alpha, \beta : \Pi_{n-1}(C) \rightarrow \mathbb{R}$ be the functions such that $\text{graph}\alpha$ and $\text{graph}\beta$ are the two cells in \mathcal{C} forming the ‘bottom’ and ‘top’ of C . Then the graph of α is compatible with d and so it is either tangent to d or transverse to d . Since $\text{graph}\alpha$ is connected, if it is tangent to d , then either $\text{graph}\alpha \subseteq L$ or $L \cap \text{graph}\alpha = \emptyset$. If the graph of α is transverse to d then by Fact 2 and the inductive hypothesis, $L \cap \text{graph}\alpha$ is definable. By

Lemma 4, $\text{fr}N \cap \text{graph}\alpha$ is a clopen subset of $L \cap \text{graph}\alpha$ and so $\text{fr}N \cap \text{graph}\alpha$ is also definable. This all also holds with the graph of β in place of the graph of α . Since N is bounded and the graph of a continuous function, $x \in \text{fr}\Pi_{n-1}(N)$ if and only if there is a y such that $(x, y) \in \text{fr}N$. So the set $\text{fr}\Pi_{n-1}(N) \cap \Pi_{n-1}(C)$ is definable. Let \mathcal{D} be a cell decomposition of $\Pi_{n-1}(C)$ compatible with $\text{fr}\Pi_{n-1}(N) \cap \Pi_{n-1}(C)$. Then for each $D \in \mathcal{D}$ we either have $D \subseteq \Pi_{n-1}(N)$ or $D \cap \Pi_{n-1}(N) = \emptyset$. For each non-open cell $D \in \mathcal{D}$ such that $D \subseteq \Pi_{n-1}(N)$, let $E_D = (\alpha|_D, \beta|_D)_D$. Take a cell decomposition of E_D compatible with d . Let E' be a cell in this decomposition such that $\text{graph}f|_D \cap E'$ is non-empty. Then by Fact 2, $\text{graph}f|_D \cap E'$ is either a finite union of Rolle leaves of the pullback of d to E' and so definable by the inductive hypothesis, or is equal to E' (in the case that E' is tangent to d). So the graph of $f|_D$ is definable. Finally, for each open cell $D \in \mathcal{D}$ such that $D \subseteq \Pi_{n-1}(N)$, the restriction of f to D is Pfaffian over $\mathcal{P}'(\mathcal{R})$ and so is definable by Lemma 3. So N is definable, as required.

□

Acknowledgements

The first author was supported in part by an EPSRC postdoctoral fellowship and by EPSRC grant EP/E050441/1, and the second author is supported by NSERC.

References

- [1] **J Bochnak, M Coste, M-F Roy**, *Real Algebraic Geometry*, Ergebnisse der Mathematik und ihrer Grenzgebiete (Series 3) 36, Springer-Verlag, Berlin, 1998; translated from the 1987 French original, revised by the authors.
- [2] **L van den Dries**, *Tame Topology and O-minimal Structures*, London Mathematical Society Lecture Note Series 248, Cambridge University Press, 1998.
- [3] **A G Khovanskii**, *Fewnomials*, Translations of Mathematical Monographs 88, American Mathematical Society, Providence, RI, 1991, translated from the Russian by Smilka Zdravkovska.
- [4] **J-M Lion, J-P Rolin**, *Volumes, feuilles de Rolle de feuilletages analytiques et théorème de Wilkie*, Ann. Fac. Sci. Toulouse Math. (6) 7 (1998), 93–112.
- [5] **J-M Lion, P Speissegger**, *The theorem of the complement for nested sub-Pfaffian sets*, Duke Math. J. 155 (2010), 35–90.
- [6] **R Moussu, C Roche**, *Théorie de Hovanskii et problème de Dulac*, Invent. Math. 105 (1991), 431–441; doi: [10.1007/BF01232274](https://doi.org/10.1007/BF01232274).

- [7] **P Speissegger**, *The Pfaffian closure of an o-minimal structure*, J. Reine Angew. Math. 508 (1999), 189–211.
- [8] **P Speissegger**, *Pfaffian sets and o-minimality*, to appear in: *Lecture Notes on O-minimal Structures and Real Analytic Geometry* (eds. C. Miller, J.-P. Rolin and P. Speissegger), Fields Institute Communications, Springer, 2012.
- [9] **A J Wilkie**, *A theorem of the complement and some new o-minimal structures*, Selecta Math. (N.S.) 5 (1999), 397–421.

Centre for Interdisciplinary Computational and Dynamical Analysis, School of Mathematics, University of Manchester, Oxford Road, Manchester, M13 9PL, UK.

Department of Mathematics & Statistics, McMaster University, 1280 Main Street West, Hamilton, Ontario L8S 4K1, Canada.

gareth.jones-3@manchester.ac.uk, speisseg@math.mcmaster.ca

Received: 17 January 2011 Revised: 25 January 2012